





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07C 237/36, A61K 31/223, A61P 35/00		A1	(11) International Publication Number: WO 00/58271
			(43) International Publication Date: 5 October 2000 (05.10.00)
(21) International Application Number: PCT/GB00/01194		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 29 March 2000 (29.03.00)			
(30) Priority Data: 9907414.8 31 March 1999 (31.03.99) GB			
(71) Applicant (for all designated States except US): CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED [GB/GB]; Cambridge House, 6-10 Cambridge Terrace, Regent's Park, London NW1 4JL (GB).			
(72) Inventors; and (75) Inventors/Applicants (for US only): SPRINGER, Caroline, Joy [GB/GB]; Institute of Cancer Research, CRC Centre for Cancer Therapeutics, 15 Cotswold Road, Sutton, Surrey SM2 5NG (GB). DAVIES, Lawrence, Christopher [GB/GB]; Institute of Cancer Research, CRC Centre for Cancer Therapeutics, 15 Cotswold Road, Sutton, Surrey SM2 5NG (GB).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(74) Agents: BRASNETT, Adrian, H. et al.; Mewburn Ellis, York House, 23 Kingsway, London WC2B 6HP (GB).			


(54) Title: NITROGEN MUSTARD COMPOUNDS AND PRODRUGS THEREFOR

(57) Abstract

This invention pertains to nitrogen mustard compounds (Formula (II)) and prodrugs thereof (Formula (I)), methods for their preparation, pharmaceutical compositions comprising such compounds, and the use of such compounds and compositions, both in vitro and in vivo, in therapy and treatment, for example, of cancer, wherein: R¹ and R² are independently -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph; R^{1a}, R^{2a}, R^{1b}, and R^{2b} are independently -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group; R³ is -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃, -CF₃, -SF₅, -SCF₃, or -CF₂CF₃; R⁴ is -H or as defined for R³; R⁵ is -H or -F; R⁷ is -H, -C(CH₃)₃, or -CH₂-CH=CH₂; Z is -CH₂-T-W; T is -CH₂-, -O-, -S-, -(S=O)-, or -(SO₂)-; W is one of: (1) -COOH; (2) -(C=O)OR⁸; (3) -(C=O)NR⁹R⁹; (4) -SO₂NHR¹⁰; (5) -SO₂OR¹¹; (6) -PO₃R¹¹R¹¹; (7) a tetrazol-5-yl group; (8) -CONH-SO₂R¹²; and, (9) -M-Het.



(I)



(II)

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

-1-

NITROGEN MUSTARD COMPOUNDS AND PRODRUGS THEREFOR

TECHNICAL FIELD

This invention pertains to nitrogen mustard compounds (Formula II) and prodrugs therefor (Formula I), methods for
5 their preparation, pharmaceutical compositions comprising such compounds, and the use of such compounds and compositions, both in vitro and in vivo, in therapy and treatment, for example, of cancer.

BACKGROUND

10 Over the years, many cytotoxic compounds have been discovered which are of potential use in cancer chemotherapy. Nitrogen mustards from one important family of such cytotoxic compounds. Nitrogen mustards are nitrogen analogs of mustard gas, $S(CH_2CH_2Cl)_2$. The clinical use of cytotoxic compounds in
15 general, and nitrogen mustards in particular, has been limited because of the poor selectivity in the cytotoxic effect between tumour cells and normal cells.

One approach to overcome this problem has involved the development of so-called prodrugs which are derivatives of
20 the cytotoxic drug, often relatively simple derivatives, whose cytotoxic properties are considerably reduced compared to those of the parent drugs. Numerous proposals have been made for the administration of such prodrugs to patients under regimes whereby the prodrug is only converted to the
25 cytotoxic drug in the region of the intended site of action, for example, by the action of an enzyme.

A variety of systems exist for delivery of the enzyme. One such system is described in WO 88/07378, and involves conjugating the enzyme to an antibody specific for a tumour

-2-


marker, delivering the antibody enzyme conjugate to a patient, allowing the conjugate to localise, and then delivering the prodrug to the patient. This system is referred to as "antibody-directed enzyme prodrug therapy"

5 (ADEPT).

Various other proposals for delivery of the enzyme to the desired site of action exist, including "gene-directed enzyme prodrug therapy" (GDEPT), in which a vector such as a viral vector carrying a gene encoding the prodrug-converting enzyme is delivered to cells at the desired site of action (Huber et al, Proc. Natl. Acad. Sci. USA, (1991) Vol. 88, p. 8039). GDEPT is also described in WO 96/03151.

A further alternative system, often referred to as "ligand-directed enzyme prodrug therapy" (LIDEPT), is to provide a ligand, generally a naturally occurring polypeptide whose biological role involves its binding to a cognate receptor on the surface of the cell, conjugated to the prodrug-activating enzyme. Such a system is described in WO 97/26918, where VEGF is particularly exemplified as an example of a ligand.

20 One class of prodrugs suggested for use in the above systems is that of prodrugs of nitrogen mustard compounds. Benzoic acid nitrogen mustards, such as the one shown below, are bifunctional alkylating agents, and a variety of prodrugs of such compounds are described in the art.




25 One class of such prodrugs comprise a protecting group which may be removed by the action of a carboxypeptidase enzyme,


-3-

such as bacterial carboxypeptidase G, including such enzymes available from Pseudomans species, such as carboxypeptidase G2 (CPG2). In many cases, the carboxylic acid group, -COOH, of the drug is protected as an amide, -CONHR, in the prodrug, for example, by reaction with an α -amino acid (e.g., glutamic acid) or a polypeptide chain, R-NH₂. The prodrug is converted into the active drug by cleavage of the amide bond, -CONH-, between the residue of the α -amino acid or oligopeptide and the residue of the benzoic acid nitrogen mustard. This enzyme, and prodrugs which may be activated by it have been described in various publications.

WO 88/07378 pertains to certain nitrogen mustard compounds of the following formula, where X and Y are -Cl and -OSO₂CH₃, or both -OSO₂CH₃, or both -OSO₂CF₃, and R is the residue of an α -amino acid RNH₂, for example, glutamic acid.




Springer et al., J. Medicinal Chem., (1990) Vol. 33, pp. 677-681 and Springer et al., Anti-Cancer Drug Design (1991), Vol. 6, pp. 467-479 pertain to certain nitrogen mustard compounds of the following formula, wherein X and Y are -Cl or -OSO₂CH₃.




- 4 -


WO 94/02450 pertains to certain nitrogen mustard compounds of the following formula, wherein X is -O-, -NH-, or -CH₂- and Y is =O.




- 5 WO 94/25429 pertains to certain 2-fluoro- and 3-fluoro-substituted nitrogen mustard compounds of the following formula, wherein the group -NH-R is the residue of an α -amino acid NH₂-R or oligopeptide NH₂-R, and Y and L, which may be the same or different in a given molecule, are leaving
10 groups:




- WO 96/03515 pertains to GDEPT systems which employ certain nitrogen mustard compounds of the following formula of the formulae shown below, in which X is -C(=O)-, -OC(=O)-, -NHC(=O)-, or -CH₂-C(=O)- and Z¹ is -O- or -NH-.



-5-



WO 96/22277 pertains to certain nitrogen mustard compounds of the following Formula, in which Z^1 is $-O-$ or $-NH-$.




- 5 Thus although a number of nitrogen mustard prodrugs exist in the art, there is a continuing need for improved prodrugs, which can provide enhanced efficacy, for example by having an improved ratio of toxicity between prodrug and active drug form, better half-life or more potency.

10


SUMMARY OF THE INVENTION

One aspect of the present invention pertains to prodrug compounds of the Formula I:



-6-

and drug compounds of the Formula II:



as described herein.

In one preferred embodiment, R¹ and R² are independently -I,
5 -Br, or -Cl.

In one preferred embodiment, R¹ and R² are both -I.

In one preferred embodiment, R^{1a}, R^{1b}, R^{2a}, R^{2b} are each
independently -H or -CH₃.

In one preferred embodiment, R^{1a}, R^{1b}, R^{2a}, R^{2b} are all -H.

10 In one preferred embodiment, R³ and R⁴ are -CF₃ and -H,
respectively.

In one preferred embodiment, R³ and R⁴ are both -F.

In one preferred embodiment, R³ and R⁴ are -CF₃ and -H,
respectively; and, R⁵ is -H.

15 In one preferred embodiment, R³ and R⁴ are both -F; and, R⁵
is -F.

In one preferred embodiment, R³ and R⁴ are both -F; and, R⁵
is -H.

In one preferred embodiment, Z is -CH₂-T-C(=O)OH or
20 -CH₂-T-C(=O)OR⁸; and, T is -CH₂-.

-7-

In one preferred embodiment, R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$.

In one preferred embodiment, the compound is selected from:

- {3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamic acid;
- {3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamic acid;
- {3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamic acid;
- {2,3,5-trifluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamic acid;
- {2,3,5-trifluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamic acid;
- {2,3,5-trifluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamic acid;
- {3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoyl}-L-glutamic acid;
- {3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamic acid; and,
- the di-tert-butyl esters thereof.

In one preferred embodiment, the compound is selected from:

- 3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoic acid;
- 3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoic acid;
- 3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoic acid;
- 2,3,5-trifluoro-4-[bis(2-chloroethyl)amino]benzoic acid;
- 2,3,5-trifluoro-4-[bis(2-bromoethyl)amino]benzoic acid;
- 2,3,5-trifluoro-4-[bis(2-iodoethyl)amino]benzoic acid;
- 3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoic acid; and
- 3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoic acid.

-8-

Another aspect of the present invention pertains to two-component systems comprising:

- (i) a first component capable of delivering a carboxypeptidase enzyme to the interior or exterior of a target cell or a vector encoding said enzyme to the interior of said cell such that said vector expresses said enzyme in said cell, and
- (ii) a prodrug as described herein capable of being converted by said enzyme into a drug as described herein.

Another aspect of the present invention pertains to kits comprising:

- (a) a prodrug compound as described herein; and,
- (b) one of:

- (i) an immunoglobulin/enzyme fusion protein or conjugate in which the immunoglobulin is specific for a cellular antigen and the enzyme is a carboxypeptidase enzyme;
- (ii) a ligand/enzyme conjugate or fusion protein, the ligand being specific for a cellular antigen and the enzyme is a carboxypeptidase enzyme;
- (iii) a vector which encodes a carboxypeptidase enzyme which can be expressed in a cell.

Another aspect of the present invention pertains to a composition comprising a drug or prodrug compound as described herein, and a pharmaceutically acceptable carrier or diluent.

Another aspect of the present invention pertains to a drug or prodrug compound as described herein for use in a method of treatment of the human or animal body.

-9-

Another aspect of the present invention pertains to a drug or prodrug compound as described herein for use in a method of treatment of cancer of the human or animal body.

5 Another aspect of the present invention pertains to use of a drug or prodrug compound as described herein for the manufacture of a medicament for use in the treatment of cancer.

10 Another aspect of the present invention pertains to a method for the treatment of cancer comprising administering to a subject suffering from cancer a therapeutically-effective amount of a drug or prodrug compound as described herein.

Another aspect of the present invention pertains to a method for the preparation of a compound according to Formula I or Formula II, as described herein.

15 BRIEF DESCRIPTION OF THE DRAWINGS


Figure 1A is a graph of survival rate versus time for Nu/Nu BalbC mice bearing a breast carcinoma xenograft treated with the compound of Example 1, {3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamic acid (denoted "A") at a
20 dosage of 600 mg/kg, as described in Example 9.

Figure 1B is a graph of survival rate versus time for Nu/Nu BalbC mice bearing a breast carcinoma xenograft treated with the prior art compound, {3-fluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamic acid (denoted "B") at a dosage of
25 1200 mg/kg, as described in Example 9.

-10-


DETAILED DESCRIPTION OF THE INVENTION

Accordingly, one aspect of the present invention pertains to a compound of general Formula I:



- 5 The compound of Formula I is a prodrug which may be converted into the corresponding active drug of Formula II by cleavage of the amide bond between the residue of R^6-NH_2 and the benzoic acid nitrogen mustard residue.

Thus, another aspect of the present invention pertains to a
10 compound of general Formula II:



In the above Formulae I and II:

R^1 is -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph ;

R^2 is -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph ;

- 15 wherein Ph denotes a phenyl group which is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from a C₁₋₄ alkyl group, -F, -Cl, -Br, -I, -CN, or -NO₂;

-11-

R^{1a} is -H, a C_{1-4} alkyl group, or a C_{1-4} haloalkyl group ;

R^{2a} is -H, a C_{1-4} alkyl group, or a C_{1-4} haloalkyl group ;

R^{1b} is -H, a C_{1-4} alkyl group, or a C_{1-4} haloalkyl group ;

R^{2b} is -H, a C_{1-4} alkyl group, or a C_{1-4} haloalkyl group ;

5 R^3 is -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃, -CF₃,
-SF₅, -SCF₃, or -CF₂CF₃ ;

R^4 is -H, -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃,
-CF₃, -SF₅, -SCF₃, or -CF₂CF₃ ;

R^5 is -H or -F;

10 with the proviso that if R^4 is -H, then R^3 is not -F;

R^7 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂ ;

Z is -CH₂-T-W ;

T is -CH₂-, -O-, -S-, -(S=O)-, or -(SO₂)- ;


the group -CH₂-T- may optionally be substituted with 1 or
15 2 substituents, denoted Q¹ and Q² respectively, on carbon,
wherein Q¹ and Q² are independently a C_{1-4} alkyl group or a
halogen; or, when Q¹ and Q² are bonded to adjacent carbon
atoms, Q¹ and Q² together may form a C_{3-4} alkylene radical
optionally substituted with 1, 2, 3 or 4 substituents
20 independently selected from C_{1-4} alkyl groups and halogens;

W is one of:

- (1) -COOH ;
(2) -(C=O)OR⁸ ;
(3) -(C=O)NR⁹R⁹ ;
25 (4) -SO₂NHR¹⁰ ;
(5) -SO₂OR¹¹ ;
(6) -PO₃R¹¹R¹¹ ;

-12-

(7) a tetrazol-5-yl group:

(8) $-\text{CONH}-\text{SO}_2\text{R}^{12}$; and,(9) $-\text{M}-\text{Het}$;

5 with the proviso that if T is $-\text{O}-$, $-\text{S}-$, $-(\text{S}=\text{O})-$, or $-(\text{SO}_2)-$, then W is not $-\text{COOH}$;

R^8 is a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{5-20} aryl group, or $-\text{CH}_2-\text{CH}=\text{CH}_2$;

10 wherein the C_{5-20} aryl group (e.g., $-\text{Ph}$) may optionally be substituted on carbon with from 1 to 4 substituents selected from $-\text{COOH}$, $-\text{OH}$, $-\text{NH}_2$, $-\text{CH}_2\text{NH}_2$, $-(\text{CH}_2)_{1-4}\text{COOH}$, tetrazol-5-yl, and $-\text{SO}_3\text{H}$;

15 R^9 is independently $-\text{H}$, a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{5-20} aryl group (e.g., $-\text{Ph}$), a C_{7-9} aralkyl group, or a C_{5-20} heteroaryl group linked to N via carbon;

wherein C_{5-20} heteroaryl is a 5 or 6 membered ring containing 1 to 3 heteroatoms selected from the group consisting of nitrogen and sulphur;

20 and wherein the C_{5-20} aryl group (e.g., $-\text{Ph}$), the C_{5-20} heteroaryl group, and aryl moiety of the C_{7-9} aralkyl group may optionally be substituted on carbon with from 1 to 4 substituents selected from $-\text{COOH}$, $-\text{OH}$, $-\text{NH}_2$, $-\text{CH}_2\text{NH}_2$, $-(\text{CH}_2)_{1-4}\text{COOH}$, tetrazol-5-yl, and $-\text{SO}_3\text{H}$;

25 and wherein the C_{3-6} cycloalkyl group may optionally carry a methyl group;

R^{10} is a C_{1-6} alkyl group, $-\text{CH}_2-\text{CH}=\text{CH}_2$, a C_{3-6} cycloalkyl group, a C_{1-4} haloalkyl group (e.g., $-\text{CF}_3$, $-\text{CH}_2\text{CF}_3$), or a C_{5-20} aryl group (e.g., phenyl);

-13-

wherein the C₅₋₂₀aryl group (e.g., -Ph), the C₅₋₂₀heteroaryl group, and aryl moiety of the C₇₋₉aralkyl group may optionally be substituted on carbon with from 1 to 4 substituents selected from -COOH, -OH, -NH₂, -CH₂NH₂,

5 -(CH₂)₁₋₄COOH, tetrazol-5-yl, and -SO₃H;

and wherein the C₃₋₆cycloalkyl group may optionally carry a methyl group;

R¹¹ represents -H, a C₁₋₆alkyl group, or a C₃₋₆cycloalkyl group;

10 R¹² is one of:

(a) a C₃₋₇cycloalkyl group;

(b) a C₁₋₆alkyl group, optionally substituted with one or more of: a phenyl group; a phenyl group with from 1 to 5 substituents selected from halogen, 15 -NO₂, -CF₃, C₁₋₄alkyl, C₁₋₄alkoxy, -NH₂, -NHCOCH₃, -CONH₂, -OCH₂COOH, -NH(C₁₋₄alkyl), -N(C₁₋₄alkyl)₂, -NHCOOC₁₋₄alkyl, -OH, -COOH, -CN and -COOC₁₋₄alkyl; a C₁₋₄alkyl group; a C₁₋₄haloalkyl group (e.g., -CF₃); or a halogen; and,

20 (c) a C₁₋₆perfluoroalkyl group;

M represents -S-, -SO-, or -SO₂- ; and,

Het represents a 5 or 6 membered heterocyclic aromatic ring linked to M via a carbon atom of the aromatic ring, said aromatic ring containing 1, 2, 3 or 4 heteroatoms selected 25 from the group consisting of O, N and S said aromatic ring optionally being substituted on carbon atoms of the ring with 1, 2, 3 or 4 substituents selected from the group consisting of -OH, -SH, -CN, -CF₃, NH₂ and halogen.

-14-

(1) In one preferred embodiment of the compounds of Formulae I and II:

R^1 and R^2 are independently -I, -Br, or -Cl; and, the remaining groups are as defined above.

5 (2) In one preferred embodiment of the compounds of Formulae I and II:

R^1 and R^2 are both -I; and, the remaining groups are as defined above.

10 (3) In one preferred embodiment of the compounds of Formulae I and II:

R^{1a} , R^{1b} , R^{2a} , R^{2b} are each independently -H or -CH₃; and, the remaining groups are as defined above.

(4) In one preferred embodiment of the compounds of Formulae I and II:

15 R^{1a} , R^{1b} , R^{2a} , R^{2b} are all -H; and, the remaining groups are as defined above.

(5) In one preferred embodiment of the compounds of Formulae I and II:

20 R^1 and R^2 are independently -I, -Br, or -Cl; R^{1a} , R^{1b} , R^{2a} , R^{2b} are each independently -H or -CH₃; and, the remaining groups are as defined above.

(6) In one preferred embodiment of the compounds of Formulae I and II:

25 R^1 and R^2 are both -I; R^{1a} , R^{1b} , R^{2a} , R^{2b} are all -H; and, the remaining groups are as defined above.

-15-

(7) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are $-CF_3$ and $-H$, respectively; and,
the remaining groups are as defined above.

5 (8) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are both $-F$; and,
the remaining groups are as defined above.

10 (9) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are $-CF_3$ and $-H$, respectively;
 R^5 is $-H$; and,
the remaining groups are as defined above.

15 (10) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are both $-F$;
 R^5 is $-F$; and,
the remaining groups are as defined above.

20 (11) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are both $-F$;
 R^5 is $-H$; and,
the remaining groups are as defined above.

25 (12) In one preferred embodiment of the compounds of Formulae I and II:

R^1 and R^2 are independently $-I$, $-Br$, or $-Cl$;
 R^{1a} , R^{1b} , R^{2a} , R^{2b} are each independently $-H$ or $-CH_3$;
 R^3 and R^4 are $-CF_3$ and $-H$, respectively, and R^5 is $-H$; or
 R^3 and R^4 are both $-F$, and R^5 is $-F$; or
30 R^3 and R^4 are both $-F$, and R^5 is $-H$; and,

-16-

the remaining groups are as defined above.

(13) In one preferred embodiment of the compounds of Formulae I and II:

R^1 and R^2 are both -I;

5 R^{1a} , R^{1b} , R^{2a} , R^{2b} are all -H;

R^3 and R^4 are both -F, and R^5 is -F; or

R^3 and R^4 are both -F, and R^5 is -H; and,

the remaining groups are as defined above.

(14) In one preferred embodiment of the compounds of

10 Formula I:

R^7 is -H, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$ and R^8 is as defined above; and,

the remaining groups are as defined above.

15 (15) In one preferred embodiment of the compounds of Formula I:

R^7 is -H or $-C(CH_3)_3$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$ and R^8 is as defined above; and,

20 the remaining groups are as defined above.

(16) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are $-CF_3$ and -H, respectively;

R^7 is -H, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

25 Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$ and R^8 is as defined above; and,

the remaining groups are as defined above.

(17) In one preferred embodiment of the compounds of Formulae I and II:

30 R^3 and R^4 are both -F;

-17-

R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$ and R^8 is as defined above; and,
the remaining groups are as defined above.

5 (18) In one preferred embodiment of the compounds of
Formulae I and II:

R^3 and R^4 are $-CF_3$ and $-H$, respectively;

R^5 is $-H$;

R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

10 Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$ and R^8 is as defined above; and,
the remaining groups are as defined above.

(19) In one preferred embodiment of the compounds of
Formulae I and II:

15 R^3 and R^4 are both $-F$;

R^5 is $-F$;

R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

20 T is $-CH_2-$ and R^8 is as defined above; and,
the remaining groups are as defined above.

(20) In one preferred embodiment of the compounds of
Formulae I and II:

R^3 and R^4 are both $-F$;

R^5 is $-H$;

25 R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$ and R^8 is as defined above; and,
the remaining groups are as defined above.

30 (21) In one preferred embodiment of the compounds of
Formulae I and II:

R^1 and R^2 are independently $-I$, $-Br$, or $-Cl$;

-18-

R^{1a} , R^{1b} , R^{2a} , R^{2b} are each independently -H or -CH₃;
 R^3 and R^4 are -CF₃ and -H, respectively, and R^5 is -H; or
 R^3 and R^4 are both -F, and R^5 is -F; or
 R^3 and R^4 are both -F, and R^5 is -H;
5 R^7 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂;
 Z is -CH₂-T-C(=O)OH or -CH₂-T-C(=O)OR⁸, wherein:
T is -CH₂- and R⁸ is as defined above.

(22) In one preferred embodiment of the compounds of
Formulae I and II:

10 R^1 and R^2 are both -I;
 R^{1a} , R^{1b} , R^{2a} , R^{2b} are all -H;
 R^3 and R^4 are both -F, and R^5 is -F; or
 R^3 and R^4 are both -F, and R^5 is -H;
 R^7 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂;
15 Z is -CH₂-T-C(=O)OH or -CH₂-T-C(=O)OR⁸, wherein:
T is -CH₂- and R⁸ is as defined above.

(23) In one preferred embodiment of the compounds of
Formula I:

R^7 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂;
20 Z is -CH₂-T-C(=O)OH or -CH₂-T-C(=O)OR⁸, wherein:
T is -CH₂- ; and,
 R^8 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂; and,
the remaining groups are as defined above.

(24) In one preferred embodiment of the compounds of

25 Formula I:
 R^7 is -H or -C(CH₃)₃;
 Z is -CH₂-T-C(=O)OH or -CH₂-T-C(=O)OR⁸, wherein:
T is -CH₂- ; and,
 R^8 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂; and,
30 the remaining groups are as defined above.

-19-

(25) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are $-CF_3$ and $-H$, respectively;

R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

5 Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$; and,

R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$; and,

the remaining groups are as defined above.

10 (26) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are both $-F$;

R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$; and,

15 R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$; and,

the remaining groups are as defined above.

(27) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are $-CF_3$ and $-H$, respectively;

20 R^5 is $-H$;

R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$; and,

R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$; and,

25 the remaining groups are as defined above.

(28) In one preferred embodiment of the compounds of Formulae I and II:

R^3 and R^4 are both $-F$;

R^5 is $-F$;

30 R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;

Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:

T is $-CH_2-$; and,

-20-

R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$; and,
the remaining groups are as defined above.

(29) In one preferred embodiment of the compounds of
Formulae I and II:

- 5 R^3 and R^4 are both $-F$;
 R^5 is $-H$;
 R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;
 Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:
 T is $-CH_2-$; and,
10 R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$; and,
the remaining groups are as defined above.

(30) In one preferred embodiment of the compounds of
Formulae I and II:

- R^1 and R^2 are independently $-I$, $-Br$, or $-Cl$;
15 R^{1a} , R^{1b} , R^{2a} , R^{2b} are each independently $-H$ or $-CH_3$;
 R^3 and R^4 are $-CF_3$ and $-H$, respectively, and R^5 is $-H$; or
 R^3 and R^4 are both $-F$, and R^5 is $-F$; or
 R^3 and R^4 are both $-F$, and R^5 is $-H$;
 R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;
20 Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:
 T is $-CH_2-$; and,
 R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$.

(31) In one preferred embodiment of the compounds of
Formulae I and II:

- 25 R^1 and R^2 are both $-I$;
 R^{1a} , R^{1b} , R^{2a} , R^{2b} are all $-H$;
 R^3 and R^4 are both $-F$, and R^5 is $-F$; or
 R^3 and R^4 are both $-F$, and R^5 is $-H$;
 R^7 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$;
30 Z is $-CH_2-T-C(=O)OH$ or $-CH_2-T-C(=O)OR^8$, wherein:
 T is $-CH_2-$; and,
 R^8 is $-H$, $-C(CH_3)_3$, or $-CH_2-CH=CH_2$.

-21-

In one preferred embodiment, the compound is {3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamic acid, as shown in Example 1(iv).

5 In one preferred embodiment, the compound is {3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamic acid, as shown in Example 2(ii).

In one preferred embodiment, the compound is {3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamic acid, as shown in Example 3(ii).

10 In one preferred embodiment, the compound is {2,3,5-trifluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamic acid, as shown in Example 4(iv).

In one preferred embodiment, the compound is {2,3,5-trifluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamic acid,
15 as shown in Example 5(ii).

In one preferred embodiment, the compound is {2,3,5-trifluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamic acid, as shown in Example 6(ii).

In one preferred embodiment, the compound is {3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoyl}-L-glutamic acid, as shown
20 in Example 7(iv).

In one preferred embodiment, the compound is {3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamic acid, as shown in Example 8(iv).

25 Compounds of Formula I in which (a) R^7 is other than hydrogen; and, (b) W is $-(C=O)-O-R^8$ where R^8 is other than hydrogen (e.g., diesters) are usually made as intermediates in the

-22-

production of compounds in which R⁷ and R⁸ are both H (e.g., diacids), which are preferred for use as prodrugs. This is so as to provide protection of the carboxy groups of the compound during synthesis of the nitrogen mustard moiety, as shown in the Examples below. Removal of such protecting groups is illustrated in the Examples, but is also described in WO 88/07378 and WO 90/02729. Examples of such compounds are discussed below.

In one preferred embodiment, the compound is di-*tert*-butyl {3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamate, as shown in Example 1(iii).

In one preferred embodiment, the compound is di-*tert*-butyl {3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamate, as shown in Example 2(i).

In one preferred embodiment, the compound is di-*tert*-butyl {3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamate, as shown in Example 3(i).

In one preferred embodiment, the compound is di-*tert*-butyl {2,3,5-trifluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamate, as shown in Example 4(iii).

In one preferred embodiment, the compound is di-*tert*-butyl {2,3,5-trifluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamate, as shown in Example 5(i).

In one preferred embodiment, the compound is di-*tert*-butyl {2,3,5-trifluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamate, as shown in Example 6(i).

-23-

In one preferred embodiment, the compound is di-*tert*-butyl (3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoyl)-L-glutamate, as shown in Example 7(iii).

5 In one preferred embodiment, the compound is di-*tert*-butyl (3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoyl)-L-glutamate, as shown in Example 8(iii).

Salts

10 It may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the compounds of Formula I and II, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge et al., 1977.

15 For example, if the compound is anionic, or has a functional group which may be anionic (e.g., -COOH may be -COO⁻), then a salt may be formed with a suitable cation. Examples of suitable inorganic cations include, but are not limited to, alkali metal ions such as Na⁺ and K⁺, alkaline earth cations such as Ca²⁺ and Mg²⁺, and other cations such as Al³⁺. Examples of suitable organic cations include, but are not limited to, ammonium ion (i.e., NH₄⁺) and substituted ammonium ions (e.g., NH₃R⁺, NH₂R₂⁺, NHR₃⁺, NR₄⁺). Examples of some suitable substituted ammonium ions are those derived from: ethylamine, diethylamine, ethylenediamine, ethanolamine, diethanolamine, piperazine. An example of a common quaternary ammonium ion 25 is N(CH₃)₄⁺.

If the compound is cationic, or has a functional group which may be cationic (e.g., -NH₂ may be -NH₃⁺), then a salt may be formed with a suitable anion. Examples of suitable inorganic anions include, but are not limited to, those derived from 30 the following inorganic acids: hydrochloric, hydrobromic,

-24-


hydroiodic, sulfuric, sulfurous, nitric, nitrous, phosphoric, and phosphorous. Examples of suitable organic anions include, but are not limited to, anions from the following organic acids: acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetyoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and valeric.

10 Methods of Synthesis

In a further aspect, the present invention provides methods for the preparation of compounds of Formulae I and II.


Compounds of Formula I in which (a) R^7 is hydrogen; and, (b) W is $-(C=O)-O-R^8$ where R^8 is hydrogen (e.g., diacids) may be prepared from the corresponding compounds, where in (a) R^7 is other than hydrogen; and, (b) W is $-(C=O)-O-R^8$ where R^8 is other than hydrogen (e.g., diesters). A suitable method is by treatment with a suitable acid, for example in a substantially non-aqueous medium, for example, at room temperature (e.g., at 15-25°C). Examples of suitable acids include trifluoroacetic acid (TFA) and formic acid (HCOOH). The process may include converting the deprotected compound (e.g., diacid) into a pharmaceutically acceptable salt thereof.

-25-



Prodrugs of the invention may thus be produced by processes analogous to those described in WO 88/07378, WO 90/02729 and WO 91/03460.

- 5 Bismesyl compounds of the invention, in which R¹ and R² are mesyl (i.e., -SO₂CH₃), R⁷ is not hydroxy, and Z does not include an unprotected hydroxy group may be made from the corresponding di-hydroxy compound by reaction with methane sulphonic anhydride ((CH₃SO₂)₂O) in CH₂Cl₂ and triethylamine
- 10 (NET₃) at room temperature.




-26-

Similarly, optionally substituted phenyl sulphonic anhydride (e.g., $(\text{PhSO}_2)_2\text{O}$) may be used under similar conditions to provide compounds of the invention in which R^1 and R^2 are optionally substituted $-\text{OSO}_2\text{Ph}$.


- 5 The protected bismesyl compounds described above may be converted into other compounds of the invention in which R^7 is $-\text{OH}$ and Z includes an unprotected hydroxy group, by deprotection in TFA as described in the examples herein.

- Alternatively, the protected bismesyl compounds may be used
 10 to provide further compounds of the invention by treatment with an alkali metal halide (e.g., LiCl , LiBr , or NaI) in a suitable solvent (e.g., acetone or N,N -dimethylacetamide) under elevated temperature or under reflux, to cause
 15 substitution of one or both mesyl groups by the corresponding halide (X denotes Cl , Br , or I). The resulting compound may be deprotected as above.




-27-

Certain compounds of the Formula I may be made by reaction of the corresponding compound of Formula II by reaction with a compound of formula $\text{H}_2\text{N}-\text{CH}(\text{Z})\text{C}(=\text{O})\text{OR}^7$, where Z and R^7 are as defined above, in a suitable solvent (e.g., DMF, N,N-dimethylformamide) in the presence of diethyl cyanophosphonate ($\text{NC}-\text{P}(=\text{O})(\text{OCH}_2\text{CH}_3)_2$) and base.




The benzoic acid compound referred to above may be prepared in any convenient way. A suitable means is by reaction of the corresponding nitrile with a strong base (e.g., NaOH) in an aqueous solvent (e.g., ethanol) under reflux.




-28-

The nitrile referred to above may be obtained by reaction of a substituted benzonitrile, where G is a group, such as F or Cl, which is capable of being substituted with (optionally substituted) diethanolamine or protected diethanolamine in a suitable solvent, such as N,N-dimethylacetamide (DMA).




Alternatively, certain compounds of Formula I, where R^3 , R^4 and R^5 are all fluoro, R^7 is other than OH , and Z does not include an unprotected OH group, may be prepared from the corresponding substituted benzamide, where G is as defined above, by reaction with (optionally substituted) diethanolamine in a suitable solvent such as N,N-dimethylacetamide.

-29-



The substituted benzamide referred to above may be obtained by reaction of the corresponding benzoyl chloride, where R^3 , R^4 and R^5 are $-F$, with a compound of formula $H_2N-CH(Z)C(=O)OR^7$, with Z and R^7 defined above.



The methods described both above and in the Examples below, alone or in combination with other known methods, may be readily adapted by one of ordinary skill in the art to obtain other compounds of the present invention.

-30-

Compositions, Systems, Kits, and Uses

- The compounds of the present invention, and particularly the prodrugs of Formula I, are suitable for use in a method of treatment of the human or animal body by therapy,
- 5 particularly a method of treatment of cancer or other conditions in which selective killing of target cells in the body of a patient is required. The animal may be a non-human mammal. The cancer may be any disease in which there is a neoplastic cell growth, including leukemias and solid
- 10 tumours. Examples of tumours which may be treated in accordance with the present invention include breast, colorectal and ovarian tumours, as well as pancreatic, melanoma, glioblastoma, hepatoma, small cell lung, non-small cell lung, muscle and prostate tumours.
- 15 It will be understood by those of skill in the art that the treatment of a tumour includes any measure taken by a physician to alleviate the effect of the tumour on a patient. Thus, although complete remission of the tumour is a desirable goal, effective treatment will also include any
- 20 measure capable of achieving partial remission of the tumour as well as a slowing down in the rate of growth of a tumour including metastases. Such measures can be effective in prolonging and/or enhancing the quality of life and/or relieving the symptoms of the disease.
- 25 The present invention thus extends in various aspects to the use of a prodrug of the invention in a method of medical treatment, particularly a method of treatment of cancer; the use of such a prodrug in the manufacture of a composition for the treatment of a disease, disorder or other medical
- 30 condition, e.g. in the manufacture of a composition for the treatment of cancer; a method of treating a human or animal patient suffering from cancer, the method comprising

-31-

administering to the patient a prodrug of the invention in combination with an enzyme-prodrug therapy; and a method of making a pharmaceutical composition comprising admixing a prodrug of the invention with a pharmaceutically acceptable
5 excipient, vehicle or carrier, and optionally other ingredients.

A prodrug of the invention may be selectively converted to the corresponding active drug by the enzyme component of an enzyme-prodrug therapy such as ADEPT, GDEPT or LIDEPT system.
10 Accordingly, the present invention provides a two component system comprising:

- (i) a first component capable of delivering a carboxypeptidase enzyme to the interior or exterior of a target cell or a vector encoding said enzyme
15 to the interior of said cell such that said vector expresses said enzyme in said cell, and
- (ii) a prodrug of Formula I capable of being converted by said enzyme into a drug of Formula II.

It will be apparent to those of skill in the art that the
20 first component may be in the form of an antibody or ligand conjugated to the enzyme, or a vector capable of expressing a carboxypeptidase enzyme, or indeed any other suitable form for delivery of the enzyme to the cell such that activation of a prodrug may occur selectively at the site of the cell.
25 Such means are well known and described as such in the prior art mentioned above.

The term "antibody" should be construed as covering antibody derivatives, functional equivalents and homologues of antibodies, including synthetic molecules and molecules whose
30 shape mimics that of an antibody enabling it to bind an antigen or epitope. Thus the term includes monoclonal and

-32-

polyclonal antibodies, as well as antibody fragments capable of binding an antigen such as Fab fragments, Fv fragments consisting of the VL and VH domains of a single arm of an antibody or single chain Fvs in which said domains are linked
5 by a polypeptide linker.

Examples of suitable ligands include vascular endothelial growth factor (VEGF), epidermal growth factor, EGF, (which binds to epidermal growth factor receptor), heregulin and c-erbB2 ligand. The known VEGF receptors are tyrosine
10 kinases called flt-1 and flk-1, which are specifically expressed in endothelial cells of the tumour and in the border between the tumour and normal tissue. EGFR and c-ErbB2 are expressed in a number of tumour types.

For GDEPT, a wide variety of vectors are available. These
15 include those which are based upon a retrovirus. Such vectors are widely available in the art. Huber et al (ibid) report the use of amphotropic retroviruses for the transformation of hepatoma, breast, colon or skin cells. Culver et al (Science (1992), Vol. 256, pp. 1550-1552) also
20 describe the use of retroviral vectors in GDEPT. Such vectors or vectors derived from such vectors may also be used. Other retroviruses may also be used to make vectors suitable for use in the present invention. Such retroviruses include rous sarcoma virus (RSV). The promoters from such
25 viruses may be used in vectors in a manner analogous to that described above for MLV.

EP-A-415 731 describes molecular chimeras comprising a promoter which may be activated in a tumour cell operably linked to a heterologous gene encoding an enzyme capable of
30 converting a prodrug into a cytotoxic agent. Such molecular chimeras may be used to express carboxypeptidase in tumour cells in order to activate prodrugs of the invention.

-33-

EP-A-415 731 describes incorporation of such molecular chimeras into viral vectors, e.g. adenoviral or retroviral vectors. Such viral vectors may also be adapted for utilization in the present invention.

- 5 Other recombinant viral vector delivery systems are described in WO 91/02805, WO 92/14829, WO 93/10814, WO 94/21792, WO 95/07994, WO 95/14091 and WO 96/22277. Methods for producing vector delivery systems based on the above-mentioned disclosures may be used to deliver vectors encoding
10 the activating enzyme (e.g. carboxypeptidase) to target cells.

- Englehardt et al. (Nature Genetics (1993), Vol. 4, pp. 27-34) describes the use of adenovirus based vectors in the delivery of the cystic fibrosis transmembrane conductance product
15 (CFTR) into cells, and such adenovirus based vectors may also be used in accordance with the present invention. Vectors utilising adenovirus promoter and other control sequences may be of use in delivering a system according to the invention to cells in the lung, and hence useful in treating lung
20 tumours.

- Vectors encoding the carboxypeptidase may be made using recombinant DNA techniques known per se in the art. The sequences encoding the enzyme may be constructed by splicing synthetic or recombinant nucleic acid sequences together, or
25 modifying existing sequences by techniques such as site directed mutagenesis. Reference may be made to "Molecular Cloning" by Sambrook et al (1989, Cold Spring Harbor) for discussion of standard recombinant DNA techniques. In general, the vector may be any DNA or RNA vector used in
30 GDEPT therapies.

-34-

The carboxypeptidase will be expressed from the vector using a promoter capable of being expressed in the cell to which the vector is targeted. The promoter will be operably linked to the sequences encoding the enzyme and its associated sequences.

Suitable promoters include viral promoters such as mammalian retrovirus or DNA virus promoters, e.g. MLV, CMV, RSV and adenovirus promoters. Preferred adenovirus promoters are the adenovirus early gene promoters. Strong mammalian promoters may also be suitable. An example of such a promoter is the EF-1 α promoter which may be obtained by reference to Mizushima and Nagata (Nucl. Acids Res. (1990), Vol. 18, p. 5322). Variants of such promoters retaining substantially similar transcriptional activities may also be used.

The c-erbB2 proto-oncogene is expressed in breast tissues at low levels and in a tissue restricted manner. In some tumour states, however, the expression of this protein is increased, due to enhanced transcriptional activity. Notable examples of this are breast tissue (about 30% of tumours), ovarian (about 20%) and pancreatic tumours (about 50-75%). In such tumours where expression of c-erbB2 is increased due to enhanced transcription or translation, the c-erbB2 promoter may be used to direct expression of the activating enzyme in a cell specific manner. The specificity of GDEPT may be increased since transfection of normal cells by a vector with a c-erbB2 promoter will provide only very limited amount of enzyme or none and thus limited activation of prodrug. The use of the c-erbB2 promoter and homologous promoters in GDEPT is more fully described in WO 96/03151.

A carboxypeptidase enzyme of the invention will be any suitable carboxypeptidase enzyme capable of cleaving the glutamic acid or glutamic acid derivative moiety of compounds

-35-

of the Formula I. Preferred carboxypeptidase enzymes are bacterial carboxypeptidases, particularly carboxypeptidase G enzymes including CPG1, 2 or 3. A pseudomonad carboxypeptidase enzyme is described in Levy and Goldman, J. Biol. Chem., 1967, 12, 2933-2938. CPG1 is described by Cornell and Charm, Biotechnology and Bioengineering, Vol XVIII, 1976, 1171-1173. CPG2 is described in WO 88/07378 and CPG3 is described by Yasuda et al, Bull. Mukogawa, Women's Univ. Nat. Sci, 1994, 42, 63-66. Although naturally occurring carboxypeptidase enzymes are preferred as a matter of convenience, synthetic variants of such enzymes may be made. For example, WO 96/03515 described synthetic variants in which one or more of the glycosylation motifs (Asn-Xaa-Ser/Thr) (where Xaa is any amino acids) are altered to prevent glycosylation. For example, within the primary amino acid sequence of CPG2, there are three such consensus motifs, located at residues Asn 222, Asn 264 and Asn 272. Alteration of one or more of these sites is preferred when CPG2 is used in the present invention. Desirably, the alteration is substitution to leucine or glutamine. Such modifications may also be made in the practice of the present invention.

Other variants of naturally occurring sequences may also be made provided such variants retain carboxypeptidase activity. For example, variants with amino acid sequences having at least 50%, preferably 60% more preferably 70%, 80%, 85%, 90% or 95% amino acid identity may be used. Amino acid identity may be determined by standard algorithms available in the art such as the BLAST algorithms using default parameters.

Where variations of the naturally occurring sequences are made these will be such that the enzyme retains its ability to convert the prodrugs of the invention to their corresponding active drugs at substantially the same rate as the unchanged, unglycosylated enzyme. In this context, "substantially

-36-

unchanged" will desirably be within 1 order of magnitude, and preferably from about 2-fold less to 2 or 5 fold more.

Carboxypeptidase enzymes may be isolated from natural sources or produced recombinantly, for example, as described in

5 WO 96/03515.

The carboxypeptidase may be expressed as a fusion with a signal sequence which directs the enzyme to the surface of a cell. Such a system is described in WO 96/03515.

10 Prodrugs of the present invention may also be used as reagents in *in vitro* systems to test the activity of candidate carboxypeptidase enzymes for incorporation into ADEPT, LIDEPT or GDEPT systems or to test cells such as biopsy samples for their suitability for treatment with an enzyme prodrug system.

15 Such model systems may also be used to test the toxicity, activity or selectivity of compounds of the present invention against a range of tumour cell types. Suitable tumour cells for use in these model systems include ovarian, colon, glioma, breast, small cell and non-small cell lung cancer
20 cells, and melanomas.

In a further aspect, the present invention provides a pharmaceutical composition, medicament, drug or other composition comprising a prodrug of the invention. The composition may include a pharmaceutically acceptable carrier
25 or diluent.

The invention also provides a kit comprising:

- (a) a prodrug of the invention; and,
- (b) one of:

-37-

- (i) an immunoglobulin/enzyme fusion protein or conjugate in which the immunoglobulin is specific for a cellular (e.g., tumour-associated) antigen and the enzyme is a carboxypeptidase enzyme;
- 5 (ii) a ligand/enzyme conjugate or fusion protein, the ligand being specific for a cellular (e.g., tumour associated) antigen and the enzyme is a carboxypeptidase enzyme;
- 10 (iii) a vector which encodes a carboxypeptidase enzyme which can be expressed in a cell (e.g. tumour cell).

In the kits of the invention, the vectors conjugates or fusion proteins may themselves be provided in a composition including a pharmaceutically acceptable carrier or diluent.

15 Optionally, the kit may include suitable instructions for its use.

Compositions according to the present invention, and for use in accordance with the present invention, may include, in addition to active ingredient, a pharmaceutically acceptable

20 excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of

25 administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous or intravenous.

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient may be in the form of a parenterally acceptable aqueous

30 solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for

-38-

example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection or Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

- 5 In ADEPT, the prodrug will normally be administered parenterally, e.g. intravenously or intraperitoneally. Thus, the pharmaceutical composition of the invention may be one which is suitable for parenteral administration. Such a composition conveniently contains the prodrug and isotonic
10 saline or bicarbonate as diluent.

GDEPT will normally involve parenteral administration of both the prodrug and the vector encoding the carboxypeptidase enzyme. Administration by the intravenous route is frequently found to be the most practical. For glioblastoma
15 the route is often intratumoural.

Liposomes containing the vector may be administered by injection in a sterile solution. Viruses may be isolated from packaging cell lines and may be administered by regional perfusion or direct intratumoural injection, or direct
20 injection into a body cavity (intracavitary administration), for example by intra-peritoneum injection.

Compositions according to the present invention, and for use in accordance with the present invention, may be presented in unit-dose or multi-dose containers, for example sealed
25 ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example water, for injections, immediately prior to use. Injection solutions and suspensions may be prepared extemporaneously from sterile
30 powders, granules and tablets of the kind previously described.

-39-

Administration of the prodrug and/or vector and/or fusion and/or conjugate is preferably in a "therapeutically effective amount", that being sufficient to show benefit to the patient.

5

The doses of each component and the route and time-course of their administration will ultimately be at the discretion of the physician, who will take into account such factors as the nature and severity of what is being treated and the age, weight and condition of the patient.

10

Suitable doses of prodrug and conjugate for the ADEPT approach are given in Bagshawe et al. Antibody, Immunoconjugates, and Radiopharmaceuticals (1991), Vol. 4, pp. 915-922. A suitable dose of conjugate may be from 2000 to 200,000 enzyme units/m² (e.g. 20,000 enzyme units/m²) and a suitable dose of prodrug may be from 20 to 2000 mg/m² (e.g. 200 mg/m²).

15

In order to secure maximum concentration of the fusion protein or conjugate at the site of desired treatment, it is normally desirable to space apart administration of the two components by at least 4 hours. The exact regime will be influenced by various factors including the nature of the tumour to be targeted and the exact nature of the prodrug. A typical regime is to administer the conjugate at 0 h, galactosylated clearing antibody at 24 h, and prodrug at 48 h. If no clearing antibody is used, it would generally be longer than 48 h before the prodrug could be injected.

20

25

In using the LIDEPT systems of the present invention the prodrug will usually be administered following administration of the ligand-enzyme fusion protein or conjugate. Typically, the ligand/enzyme will be administered to the patient, and its uptake monitored, for example by recovery and analysis of

30

-40-

a biopsy sample of targeted tissue or by injecting trace-labelled protein ligand enzyme.

In using the GDEPT system the prodrug may be administered following administration of the vector encoding the

- 5 activating enzyme. Typically, the vector will be administered to the patient and then the uptake of the vector by transfected or infected (in the case of viral vectors) cells monitored, for example by recovery and analysis of a biopsy sample of targeted tissue.
- 10 The amount of vector delivered will be such as to provide an effective cellular concentration of enzyme so that the prodrug may be activated in sufficient concentration at the site of a tumour to achieve a therapeutic effect, e.g. reduction in the tumour size. This may be determined by
- 15 clinical trials which involve administering a range of trial doses to a patient and measuring the degree of infection or transfection of a target cell or tumour. The amount of prodrug required will be similar to or greater than that for ADEPT systems of the type mentioned above.
- 20 A treatment according to the present invention may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.


EXAMPLES

- 25 The following are examples are provided solely to illustrate the present invention and are not intended to limit the scope of the invention, as described herein.

-41-


Example 1

(i) 3,5-difluoro-4-[bis(2-hydroxyethyl)amino]benzonitrile



A solution of 3,4,5-trifluorobenzonitrile (5 g, 32 mmol) and diethanolamine (8 ml, 85 mmol) in N,N-dimethylacetamide was stirred for 10 days. The solvent was then evaporated, the residue partitioned between CH_2Cl_2 (200 ml) and H_2O (200 ml), the organic layer dried (MgSO_4) and evaporated to dryness. The residue was chromatographed using CH_2Cl_2 -EtOAc as eluent to give, after recrystallization from toluene, 2.7 g (35%) of pure white crystals: mp 63-65°C; ^1H NMR δ 3.33 (t, 4H, NCH_2 , $J = 6\text{Hz}$), 3.48 (t, 4H, OCH_2 , $J = 6\text{Hz}$), 4.43 (bs, 2H, OH), 7.57 (dd, 2H, H2+6, $J_{\text{H}2, \text{F}3 \text{ and } \text{H}6, \text{F}5} = 8.5\text{Hz}$, $J_{\text{H}2, \text{F}5 \text{ and } \text{H}6, \text{F}3} = 2\text{Hz} = 2\text{Hz}$); ^{19}F NMR δ -116.0 (d, 2F, $\text{F}3+5$, $J_{\text{F}3, \text{H}2 \text{ and } \text{F}5, 6} = 7.5\text{Hz}$); MS m/z 265 ($\text{M}+\text{Na}^+$, 10), 243 ($\text{M}+\text{H}^+$, 100), 211 ($\text{M}-\text{CH}_2\text{OH}$, 85). Anal. ($\text{C}_{11}\text{H}_{12}\text{N}_2\text{O}_2\text{F}_2$) C, H, N.

(ii) 3,5-difluoro-4-[bis(2-hydroxyethyl)amino]benzoic acid




A solution of the 3,5-Difluoro-4-[bis(2-hydroxyethyl)amino]benzonitrile (3.15 g, 13 mmol) and NaOH (5.2 g, 0.13 mol) in aqueous ethanol (65 ml, 50%) was refluxed for 2.5 hours. The solution was then partitioned between EtOAc (800 ml) and HCl (330 ml, 0.4 M), the aqueous layer washed

-42-

with EtOAc (3 x 150ml), the combined organic layers dried (MgSO₄) and evaporated to dryness. The solid recrystallized from EtOAc (100 ml) to give 2.94 g (87%) of pure white crystals: mp 146.5-150°C; ¹H NMR δ 3.3 (t, 4H, CH₂N), 3.46 (q, 4H, CH₂O, J = 5.5Hz), 4.49 (t, 2H, OH, J = 5Hz), 7.44 (d, 2H, H₂+6, J = 10.5Hz), 13.1 (bs, 1H, CO₂H); ¹⁹F NMR δ -117.3 (d, 2F, F3+5, J = 9.5Hz); MS m/z 284 (M+Na⁺, 15), 262 (M+H⁺, 100), 230 (M-CH₂OH, 40). Anal. (C₁₁H₁₃NO₄F₂) C, H, N.

(iii) di-tert-butyl {3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-glutamate



a) To a solution of di-tert-butyl-L-glutamate hydrochloride (2.66 g, 9.0 mmol) in dry dimethylformamide (135 ml) was added Et₃N (2.5 ml, 18 mmol), the 3,5-Difluoro-4-[bis(2-hydroxyethyl)amino]benzoic acid (2.35 g, 9.0 mmol) followed by diethylcyanophosphonate (1.5 ml, 9.9 ml). After stirring for 3 days the solvent was evaporated and the residue partitioned between EtOAc (450 ml) and H₂O (375 ml). The organic layer was washed with citric acid (180 ml, 10%), sat. sodium bicarbonate solution (180 ml), dried (MgSO₄) and evaporated to dryness, giving 5.6 g of the bis-hydroxy compound.


b) To a solution of the bis-hydroxy compound in dry CH₂Cl₂ (165 ml) was added 4-dimethylaminopyridine (0.22 g, 1.8 mmol) and Et₃N (6.3 ml, 45 mmol). This solution was cooled in ice, methane sulphonic anhydride (6.3 g, 36 mmol) dissolved in dry CH₂Cl₂ was added over a few minutes and the reaction was

-43-

allowed to warm up to room temperature. After 16 hr CH_2Cl_2 (150 ml) was added, the solution extracted with 10% aqueous citric acid (375 ml), dried (MgSO_4) and evaporated to dryness to give the bis-mesyl compound as a brown oil.

- 5 c) A solution of the bis-mesyl compound and NaI (13.5 g, 0.09 mol) in acetone (150 ml) was refluxed for 5 hr. The solvent was removed by evaporation, the residue partitioned between CH_2Cl_2 (375 ml) and H_2O (375 ml), the organic layer dried (MgSO_4) and evaporated to dryness. The impure product
 10 was purified on silica using CH_2Cl_2 as eluent giving 5.6 g (86%) of pure bis-iodo ester. ^1H NMR δ 1.39 + 1.41 (2s, 18H, tert-Bu), 2.0 (m, 2H, CH_2CH), 2.33 (t, 2H, CH_2CO), 3.3 (t, 4H, CH_2I), 3.57 (t, 4H, CH_2N , $J = 7\text{Hz}$), 4.3 (m, 1H, CH), 7.6 (d, 2H, H2+6, $J = 10\text{Hz}$), 8.60 (d, 1H, NH, $J = 7.5\text{Hz}$); ^{19}F NMR δ
 15 -117.2 (d, 2F, F3+5, $J = 10\text{Hz}$).

(iv) {3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-
 L-glutamic acid



- The bis-iodo ester (5.6 g, 7.8 mmol) was dissolved in
 20 trifluoroacetic acid (TFA) (140 ml). After one hour the solvent was removed by evaporation and the residue evaporated six times with EtOAc to remove TFA. The oil was re-dissolved in EtOAc, toluene was added and the solvent partly evaporated. The acid crystallized as a pure white solid
 25 (4.5 g, 95%): mp 121-123°C; ^1H NMR δ 1.95 + 2.05 (2m, 2H, CH_2CH), 2.34 (t, 2H, CH_2CO , $J = 7.5\text{Hz}$), 3.3 (t, 4H, CH_2I), 3.57 (t, 4H, CH_2N , $J = 7\text{Hz}$), 4.4 (m, 1H, CH), 7.6 (d, 2H,


-44-

H₂+6, J = 10Hz), 8.6 (d, 1H, NH, J = 7.5Hz), 12.5 (bs, 1H, CO₂H); ¹⁹F NMR δ -117.2 (d, 2F, F3+5, J = 10Hz); MS m/z 633 (M+Na⁺, 23), 611 (M+H⁺, 40), 464 (M-glu, 82). Anal. (C₁₆H₁₈N₂O₅F₂I₂) C, H, N.

5


Example 2

(i) di-tert-butyl (3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoyl)-L-glutamate



The bis-mesyl compound of Example 1(iii)(c), 2 mmol, was dissolved in N,N-dimethyl acetamide (35 ml) containing LiCl (0.85 g, 20 mmol) and heated at 85°C for 45 minutes. The solvent was removed by evaporation and the residue partitioned between CH₂Cl₂ (100 ml) and H₂O (30 ml). The organic layer was dried (MgSO₄), evaporated to dryness and the residue chromatographed using CH₂Cl₂ as eluent to give 1.11 g of product.

(ii) (3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoyl)-L-glutamic acid



-45-


The product of Example 2(i) was dissolved in 28 ml TFA and treated as in Example 1(iv) to provide 0.88 g of crystalline product.

^1H NMR δ 1.95 + 2.05 (2m, 2H, CH_2CH), 2.35 (t, 2H, CH_2CO_2 , $J = 7.5\text{Hz}$), 3.57 (t, 4H, CH_2N , $J = 6\text{Hz}$), 3.68 (t, 4H, CH_2Cl , $J = 6\text{Hz}$), 4.4 (m, 1H, CH), 7.60 (d, 2H, H2+6, $J = 10\text{Hz}$), 8.61 (d, 1H, NH, $J = 7.5\text{Hz}$); ^{19}F NMR δ -117.44 (d, 2F, F3+5, $J = 10\text{Hz}$); MS m/z 449 ($\text{M}+\text{Na}^+$, 17), 427 ($\text{M}+\text{H}^+$, 51), 377 ($\text{M}-\text{CH}_2\text{Cl}$, 42), 280 ($\text{M}-\text{glu}$, 100). Anal. ($\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}_5\text{F}_2\text{Cl}_2 \cdot 0.5\text{toluene}$) C, H, N.

10

Example 3


(i) di-tert-butyl {3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamate



The bis-mesyl compound of Example 1(iii)(c), 0.9 mmol, was dissolved in acetone (15 ml) containing LiBr (0.75 g, 9 mmol) and refluxed for 5 hours. The solvent was removed by evaporation and the residue partitioned between CH_2Cl_2 (30 ml) and H_2O (15 ml). The organic layer was dried (MgSO_4), and evaporated to dryness. The residue was chromatographed using CH_2Cl_2 -EtOAc as eluent to give 0.4 g of product.

-46-

(ii) {3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoyl}-
L-glutamic acid




The product of Example 3(i) was dissolved in 10 ml TFA and
5 treated as in Example 1(iv) to provide 0.32 g of crystalline
product.

^1H NMR δ 1.95 + 2.05 (2m, 2H, CH_2CH), 2.35 (t, 2H, CH_2CO_2 , $J = 7.5\text{Hz}$), 3.55 (t, 4H, CH_2N , $J = 5.5\text{Hz}$), 3.61 (t, 4H, CH_2Br , $J = 5.5\text{Hz}$), 4.4 (1H, m, CH), 7.60 (d, 2H, H2+6, $J = 10\text{Hz}$), 8.62
10 (d, 1H, NH, $J = 7.5\text{Hz}$); ^{19}F NMR δ -117.32 (d, 2F, F3+5, $J = 10\text{Hz}$); MS m/z 539 ($\text{M}+\text{Na}^+$, 24), 517 ($\text{M}+\text{H}^+$, 59), 423 ($\text{M}-\text{CH}_2\text{Br}$, 33), 370 ($\text{M}-\text{glu}$, 100). Anal. ($\text{C}_{16}\text{H}_{18}\text{N}_2\text{O}_5\text{F}_3\text{Cl}_2 \cdot 0.32\text{toluene}$) C, H, N.

Example 4

15 (i) di-tert-butyl (2,3,4,5-tetrafluorobenzoyl)-L-glutamate




To an ice-cold solution of di-tert-butyl L-glutamate
hydrochloride (6.6 g, 22.4 mmol) and Et_3N (6.9 ml, 50 mmol) in
dry CH_2Cl_2 (70 ml) was added, over a period of 1.5 hr,
20 2,3,4,5-tetrafluorobenzoyl chloride (5.0 g, 23.5 mmol) in dry

-47-

CH₂Cl₂ (20 ml) and thirty minutes later the solution was allowed to warm to room temperature. Sixteen hours later the solution was washed with H₂O, dried (MgSO₄) and evaporated to dryness. ¹H NMR δ 1.40 + 1.42 (2s, 18H, t-butyl), 1.85 + 1.95 (2m, 2H, CH₂CH), 2.34 (t, 2H, CH₂CO, J = 7.5Hz), 4.3 (m, 1H, CH), 7.57 (m, 1H, H₆), 8.80 (d, 1H, NH, J = 7.5Hz); ¹⁹F NMR δ -138.6 + 139.0 (2m, 2F, F₂+5), -152.4 (m, 1H, F₄), -154.9 (t, 1F, F₃, J = 20Hz).

(ii) di-tert-butyl {4-[bis(2-hydroxyethyl)amino]-2,3,5-trifluorobenzoyl}-L-glutamate




A solution of the tetrafluoroamide (4.35 g, 10 mmol) and diethanolamine (3.0 ml, 30 mmol) in N,N-dimethylacetamide (50 ml) was heated at 130°C with stirring for 2.5 days. The solvent was removed by evaporation and the residue partitioned between CH₂Cl₂ (150 ml) and H₂O (150 ml). The organic layer was dried (MgSO₄), evaporated to dryness and chromatographed using CH₂Cl₂-EtOAc as eluent to give 2.7 g (52%) of product.

¹H NMR 1.39 + 1.41 (2s, 18H, t-butyl), 1.95 (m, 2H, CH₂CH), 2.32 (t, 2H, CH₂CO₂, J = 7.5Hz), 3.3 (t, 4H, CH₂N), 3.49 (q, 4H, CH₂O, J = 5.5Hz), 4.3 (m, 1H, CH), 4.52 (t, 2H, OH, J = 5Hz), 7.18 (q, 1H, H-6), 8.48 (d, 1H, NH, J = 7.5Hz); ¹⁹F NMR δ -122.7 (t, 1F, F₅), -142.0 (m, 2F, F₂+3).

-48-

(iii) di-tert-butyl {4-[bis(2-chloroethyl)amino]-
2,3,5-trifluorobenzoyl}-L-glutamate




a) A solution of the bis-hydroxy compound (2.6 g, 5.0 mmol),
5 4-dimethylaminopyridine (90 mg, 0.7 mmol) and Et_3N (2.5 ml,
17.5 mmol) in dry CH_2Cl_2 (70 ml) was cooled in ice. Methane
sulphonic anhydride (2.5 g, 14 mmol), dissolved in dry CH_2Cl_2
(20 ml), was added over two minutes and the solution was
allowed to warm up to room temperature. After 16 hr CH_2Cl_2
10 (100 ml) was added and the solution was extracted with 10%
aqueous citric acid, dried (MgSO_4) and evaporated to dryness
to give the bis-mesyl compound as an oil (3.8 g).

b) The bis-mesyl compound was dissolved in N,N-
dimethylacetamide (25 ml) containing LiCl (0.5 g, 10 mmol).
15 After 16 hr the solvent was removed by evaporation and the
residue partitioned between CH_2Cl_2 (50 ml) and H_2O (20 ml).
The organic layer was dried (MgSO_4), evaporated to dryness and
the residue chromatographed using CH_2Cl_2 as eluent to give
1.35 g (97%) of the bis-chloro ester, as a colourless oil.

-49-


(iv) 4-[bis(2-chloroethyl)amino]-2,3,5-trifluorobenzoyl-L-glutamic acid



The bis-chloro ester (1.27 g, 2.3 mmol) was dissolved in TFA
 5 (30 ml). After one hour the solvent was evaporated and the
 residue evaporated five times with EtOAc. The product was
 re-dissolved in EtOAc, toluene was added and the solution
 evaporated to low volume. More toluene was added and the
 solution partially evaporated again giving a large mass of
 10 white crystals (1.015 g, 97.5%): mp 110°C; ^1H NMR δ 1.9 + 2.05
 (2m, 2H, CH_2CH), 2.34 (t, 2H, CH_2CO , $J = 7.5\text{Hz}$), 3.60 (t, 4H,
 CH_2N , $J = 6\text{Hz}$), 3.71 (t, 4H, CH_2Cl , $J = 6\text{Hz}$), 4.4 (m, 1H, CH),
 7.25 (m, 1H, H6), 8.56 (d, 1H, NH, $J = 8\text{Hz}$); ^{19}F NMR δ -122.67
 (t, 1F, F5, $J_{\text{F2}, 5} = 13.5\text{Hz}$), -141.3 (m, 1F, F2), -141.5 (d, 1F,
 15 F3, $J_{\text{F2}, 3} = 19\text{Hz}$); MS m/z 467 ($\text{M}+\text{Na}^+$, 40), 445 ($\text{M}+\text{H}^+$, 82), 395
 ($\text{M}-\text{CH}_2\text{Cl}$, 41), 298 ($\text{M}-\text{glu}$, 100). Anal.
 ($\text{C}_{18}\text{H}_{17}\text{N}_2\text{O}_5\text{F}_3\text{Cl}_2 \cdot 0.08\text{toluene}$) C, H, N.

Example 5


(i) di-*tert*-butyl {4-[bis(2-bromoethyl)amino]-
 20 2,3,5-trifluorobenzoyl}-L-glutamate



-50-

The bis-mesyl compound of Example 4(iii)(b), 3 mmol, was dissolved in acetone (25 ml) containing LiBr (0.87 g, 10 mmol) and refluxed for 5 hours. The solvent was removed by evaporation and the residue partitioned between CH₂Cl₂ (50 ml) and H₂O (20 ml). The organic layer was dried (MgSO₄), evaporated to dryness and the residue chromatographed using CH₂Cl₂ as eluent to give 1.33 g of product.

(ii) {4-[bis(2-bromoethyl)amino]-
2,3,5-trifluorobenzoyl}-L-glutamic acid




The product of Example 5(i) was dissolved in 30 ml TFA and treated as in Example 4(iv) to provide 1.0 g of crystalline product.

¹H NMR δ 1.90 + 2.05 (2m, 2H, CH₂CH), 2.35 (t, 2H, CH₂CO₂, J = 7.5Hz), 3.58 (t, 2H, CH₂N, J = 5.5Hz), 3.66 (t, 4H, CH₂Br, J = 5.5Hz), 4.4 (m, 1H, CH), 7.25 (dd, 1H, H₆, J_{H5, F5} = 12Hz, J_{H6, F3} = 5.5Hz), 8.56 (d, 1H, NH, J = 7.5Hz); ¹⁹F NMR δ -122.4 (m, 1F, F5), -141.5 (m, 2F, F2+3); MS m/z 557 (M+Na⁺, 31), 535 (M+H⁺, 70), 455 (M-Br, 14), 441 (M-CH₂Br, 30), 388 (M-glu, 100). Anal. (C₁₆H₁₇N₂O₅F₃Br₂·0.06toluene) C, H, N.

-51-


Example 6

(i) di-*tert*-butyl {4-[bis(2-iodoethyl)amino]-2,3,5-trifluorobenzoyl}-L-glutamate



- 5 The bis-mesyl compound of Example 4(iii)(b), 1 mmol, was dissolved in acetone (15 ml) containing NaI (1.5 g, 10 mmol) and refluxed for 5 hours. The solvent was removed by evaporation and the residue partitioned between CH₂Cl₂ (45 ml) and H₂O (15 ml). The organic layer was dried (MgSO₄),
- 10 evaporated to dryness and the residue chromatographed on silica using CH₂Cl₂ as eluent to give 0.7 g of product.

(ii) {4-[bis(2-iodoethyl)amino]-2,3,5-trifluorobenzoyl}-L-glutamic acid



- 15 The product of Example 6(i) was dissolved in 17 ml TFA and treated as in Example 4(iv) to provide 0.53 g of crystalline product.

- ¹H NMR δ 1.9 + 2.05 (2m, 2H, CH₂CH), 2.34 (t, 2H, CH₂CO₂, J = 7.5Hz), 3.3 (t, 4H, CH₂I), 3.59 (t, 4H, CH₂N, J = 7Hz), 4.38
- 20 (m, 1H, CH), 7.28 (dd, 1H, H₆, J_{H₆, F₅} = 12Hz, J_{H₆, F₃} = 2.5Hz),


-52-

8.52 (d, 1H, NH, $J = 7.5\text{Hz}$); ^{19}F NMR -122.7 (t; 1F, F5, $J_{\text{F5}, \text{H6}} = 14\text{Hz}$), -141.3 (m, 2F, F2+3); MS m/z 651 ($\text{M}+\text{Na}^+$, 23), 629 ($\text{M}+\text{H}^+$, 65), 482 ($\text{M}-\text{glu}$, 100). Anal. ($\text{C}_{16}\text{H}_{17}\text{N}_2\text{O}_5\text{F}_3\text{I}_2 \cdot 0.05\text{toluene}$) C, H, N.

5


Example 7

(i) 3,5-difluoro-4-[bis(2-hydroxypropyl)amino]benzonitrile



A solution of 3,4,5-trifluorobenzonitrile (6.3 g, 40 mmol) and diisopropanolamine (13.3 g, 0.1 mol) in N,N-
 10 dimethylacetamide (140 ml) was stirred for 3 days at 70°C. The solvent was evaporated, the residue partitioned between CH_2Cl_2 (250 ml) and H_2O (250 ml), the organic layer dried (MgSO_4) and evaporated to dryness. The residue was chromatographed using CH_2Cl_2 -EtOH as eluent.

15 (ii) 3,5-difluoro-4-[bis(2-hydroxypropyl)amino]benzoic acid




A solution of the 3,5-difluoro-4-[bis(2-hydroxypropyl)amino] benzonitrile and NaOH (8 g, 0.2 mol) in aqueous ethanol (100 ml, 50%) was refluxed for 2.5 hours. The solution was
 20 partitioned between EtOAc (1.2 l) and HCl (500 ml, 0.4 M), the aqueous layer washed with EtOAc (3 x 240 ml), the combined organic layers dried (MgSO_4) and evaporated to

-53-

dryness. The residue was chromatographed using CH_2Cl_2 -EtOH-AcOH as eluent.

(iii) di-tert-butyl {3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoyl}-L-glutamate




a) To a solution of di-tert-butyl-L-glutamate hydrochloride (0.96 g) in dry dimethylformamide (50 ml) was added Et_3N (0.9 ml, 6.4 mmol), the 3,5-difluoro-4-[bis(2-hydroxypropyl)amino]benzoic acid (0.93 g, 3.2 mmol) followed by diethyl cyanophosphonate (0.5 ml, 3.5 mmol). After 2 days the solvent was evaporated and the residue partitioned between EtOAc (150 ml) and H_2O (125 ml). The organic layer was washed with citric acid (60 ml, 10%), saturated sodium bicarbonate solution (60 ml), dried (MgSO_4), and evaporated to dryness, giving the bis-hydroxy compound.

b) To a solution of the bis-hydroxy compound in dry CH_2Cl_2 (55 ml) was added 4-dimethylaminopyridine (73 mg, 0.6 mmol) and Et_3N (2.1 ml, 15 mmol). This solution was cooled in ice, methane sulphonc anhydride (2.1 g, 12 mmol) dissolved in dry CH_2Cl_2 (18 ml) was added over a few minutes and the reaction mixture was allowed to warm up to room temperature. After a few hours CH_2Cl_2 (50 ml) was added, the solution extracted with 10% aqueous citric acid (125 ml), dried (MgSO_4), and evaporated to dryness to give the bis-mesyl compound as an oil.

-54-

c) A solution of the bis-mesyl compound and LiBr (2.61 g, 30 mmol), in acetone (50 ml) was refluxed for 5 hr. The solvent was removed by evaporation, the residue partitioned between CH₂Cl₂ (125 ml) and H₂O (125 ml), the organic layer dried (MgSO₄), and evaporated to dryness. The impure product was purified on silica using CH₂Cl₂ as eluent, to give the bis-bromo ester.

(iv) {3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoyl}-L-glutamic acid




10

The bis-bromo ester was dissolved in TFA (15 ml). After 50 minutes, the solvent was removed by evaporation and the residue evaporated four times with EtOAc to remove TFA. The oil was re-dissolved in EtOAc, toluene was added and the solvent partly evaporated. The acid crystallized as a pure white solid.

15

Example 8

(i) 3-trifluoromethyl-4-[bis(2-hydroxyethyl)amino]benzonitrile




20

-55-

A solution of 3-trifluoromethyl 4-fluoro-benzonitrile (9.45 g, 50 mmol) and diethanolamine (15.8 g, 0.15 mol) in N,N-dimethylacetamide (200 ml) was stirred for 7 days at 50°C. The solvent was evaporated, the residue partitioned between
 5 CH₂Cl₂ (400 ml) and H₂O (400 ml), the organic layer dried (MgSO₄) and evaporated to dryness. The residue was chromatographed using CH₂Cl₂-EtOH as eluent.

(ii) 3-trifluoromethyl-4-[bis(2-hydroxyethyl)amino]benzoic acid

10




15

A solution of the 3-trifluoromethyl-4-[bis(2-hydroxyethyl)amino]benzonitrile and NaOH (0.8 g, 0.02 mol) in aqueous ethanol (10 ml, 50%) was refluxed for 2 hours. The solution was partitioned between EtOAc (120 ml) and HCl (50 ml, 0.4
 15 M), the aqueous layer washed with EtOAc (3 x 25 ml), the combined organic layers dried (MgSO₄) and evaporated to dryness. The residue was recrystallized from EtOAc to give 0.39 g of white crystals.

20

(iii) di-tert-butyl {3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoyl}-L-glutamate



-56-


a) To a solution of di-tert-butyl-L-glutamate hydrochloride (0.3 g) in dry dimethylformamide (15 ml) was added Et₃N (0.28 ml, 2 mmol), the 3-trifluoromethyl-4-[bis(2-hydroxyethyl)amino]benzoic acid (0.29 g, 1 mmol) followed by diethyl cyanophosphonate (0.17 ml, 1.1 mmol). After 2 days the solvent was evaporated and the residue partitioned between EtOAc (50 ml) and H₂O (40 ml). The organic layer was washed with citric acid (20 ml, 10%), saturated sodium bicarbonate solution (20 ml), dried (MgSO₄), and evaporated to dryness, giving the bis-hydroxy compound.

b) To a solution of the bis-hydroxy compound in dry CH₂Cl₂ (6 ml) was added 4-dimethylaminopyridine (7 mg, 0.06 mmol) and Et₃N (0.21 ml, 1.5 mmol). This solution was cooled in ice, methane sulphonic anhydride (0.21 g, 1.2 mmol) dissolved in dry CH₂Cl₂ (2 ml) was added over a few minutes and the reaction mixture was allowed to warm up to room temperature. After a few hours CH₂Cl₂ (5 ml) was added, the solution extracted with 10% aqueous citric acid (12 ml), dried (MgSO₄), and evaporated to dryness to give the bis-mesyl compound as an oil.

c) A solution of the bis-mesyl compound and LiBr (0.26 g, 3 mmol) in acetone (50 ml) was refluxed for 2 hr. The solvent was removed by evaporation, the residue partitioned between CH₂Cl₂ (15 ml) and H₂O (10 ml), the organic layer dried (MgSO₄), and evaporated to dryness. The impure product was purified on silica using CH₂Cl₂ as eluent, to give the bis-bromo ester.

-57-

(iv) {3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoyl}-
L-glutamic acid



The bis-bromo ester was dissolved in TFA (6 ml). After 50
5 minutes, the solvent was removed by evaporation and the
residue washed four times with EtOAc to remove TFA. The oil
was re-dissolved in EtOAc, toluene was added and the solvent
partly evaporated. The acid crystallized as a pure white
solid (45 mg).

10

Example 9

The compound of Example 1, {3,5-difluoro-4-[bis(2-
iodoethyl)amino]benzoyl}-L-glutamic acid (denoted "A"), was
assessed in a Nu/Nu BalbC mouse bearing a breast carcinoma
xenograft that comprised 50% cells (MDA MB361) stably
15 expressing β -galactosidase plus 50% cells (MDA MB361) stably
expressing a surface tethered carboxypeptidase G2 (CPG2)
mutant enzyme (MDA MB361-STCPG2(Q)3) (see WO96/03515). This
was compared to the prior art compound, {3-fluoro-4-[bis(2-
chloroethyl)amino]benzoyl}-L-glutamic acid (denoted "B").

20

The controls comprised mice bearing the same tumour
xenograft, that were not treated with prodrug.

The results are shown in Figures 1A and 1B. As is seen,
treatment with 600 mg/kg of the compound of Example 1 ("A",
Figure 1A) provided a 50% survival rate for over 200 days,
25 whereas the untreated control group did not survive for more
than 40 days.

-58-


5 With the prior art compound ("B", Figure 1B) the survival rate of an untreated control group was comparable to the untreated group shown in Figure 1A, and although treatment provided a measurable effect, and 50% survival was found up to about 70 days post-treatment, it should be noted that to achieve this a higher dose (1200 mg/kg) of the prodrug was required. This translates on a molar basis to 0.98 mM for Example 1 (iv) and 2.9 mM for the prior art compound.

10 A number of patents and publications are cited above in order to more fully describe and disclose the invention and the state of the art to which the invention pertains. Each of these references is incorporated herein by reference in its entirety into the present disclosure.

-59-

CLAIMS

1. A compound of Formula I:



wherein:

- 5 R^1 is -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph ;
 R^2 is -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph ;
 wherein Ph denotes a phenyl group which is
 optionally substituted with 1, 2, 3, 4 or 5 substituents
 independently selected from a C₁₋₄ alkyl group, -F, -Cl,
 10 -Br, -I, -CN, or -NO₂;
- R^{1a} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;
 R^{2a} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;
 R^{1b} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;
 R^{2b} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;
- 15 R^3 is -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃, -CF₃,
 -SF₅, -SCF₃, or -CF₂CF₃ ;
- R^4 is -H, -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃,
 -CF₃, -SF₅, -SCF₃, or -CF₂CF₃ ;
- 20 R^5 is -H or -F;
- with the proviso that if R^4 is -H, then R^3 is not -F;
- R^7 is -H, -C(CH₃)₃, or -CH₂-CH=CH₂ ;

-60-


Z is $-\text{CH}_2-\text{T}-\text{W}$;

T is $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-(\text{S}=\text{O})-$, or $-(\text{SO}_2)-$;

the group $-\text{CH}_2-\text{T}-$ may optionally be substituted with 1 or 2 substituents, denoted Q^1 and Q^2 respectively, on carbon, wherein Q^1 and Q^2 are independently a C_{1-4} alkyl group or a halogen; or, when Q^1 and Q^2 are bonded to adjacent carbon atoms, Q^1 and Q^2 together may form a C_{3-4} alkylene radical optionally substituted with 1, 2, 3 or 4 substituents independently selected from C_{1-4} alkyl groups and halogens;

W is one of:

- (1) $-\text{COOH}$;
- (2) $-(\text{C}=\text{O})\text{OR}^8$;
- (3) $-(\text{C}=\text{O})\text{NR}^9\text{R}^9$;
- (4) $-\text{SO}_2\text{NHR}^{10}$;
- (5) $-\text{SO}_2\text{OR}^{11}$;
- (6) $-\text{PO}_3\text{R}^{11}\text{R}^{11}$;
- (7) a tetrazol-5-yl group:



- (8) $-\text{CONH}-\text{SO}_2\text{R}^{12}$; and,
- (9) $-\text{M}-\text{Het}$;

with the proviso that if T is $-\text{O}-$, $-\text{S}-$, $-(\text{S}=\text{O})-$, or $-(\text{SO}_2)-$, then W is not $-\text{COOH}$;

wherein:

R^8 is a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{5-20} aryl group, or $-\text{CH}_2-\text{CH}=\text{CH}_2$;

wherein the C_{5-20} aryl group may optionally be substituted on carbon with from 1 to 4 substituents

-61-

selected from -COOH, -OH, -NH₂, -CH₂NH₂, -(CH₂)₁₋₄COOH, tetrazol-5-yl, and -SO₃H;

R⁹ is independently -H, a C₁₋₆alkyl group, a C₃₋₆cycloalkyl group, a C₅₋₂₀aryl group, a C₇₋₉aralkyl group, or a C₅₋₂₀heteroaryl group linked to N via carbon;

wherein the C₅₋₂₀aryl group, the C₅₋₂₀heteroaryl group, and aryl moiety of the C₇₋₉aralkyl group may optionally be substituted on carbon with from 1 to 4 substituents selected from -COOH, -OH, -NH₂, -CH₂NH₂, -(CH₂)₁₋₄COOH, tetrazol-5-yl, and -SO₃H;

and wherein the C₃₋₆cycloalkyl group may optionally carry a methyl group;

R¹⁰ is a C₁₋₆alkyl group, -CH₂-CH=CH₂, a C₃₋₆cycloalkyl group, a C₁₋₄haloalkyl group (e.g., -CF₃, -CH₂CF₃), or a C₅₋₂₀aryl group;

wherein the C₅₋₂₀aryl group, the C₅₋₂₀heteroaryl group, and aryl moiety of the C₇₋₉aralkyl group may optionally be substituted on carbon with from 1 to 4 substituents selected from -COOH, -OH, -NH₂, -CH₂NH₂, -(CH₂)₁₋₄COOH, tetrazol-5-yl, and -SO₃H;

and wherein the C₃₋₆cycloalkyl group may optionally carry a methyl group;

R¹¹ represents -H, a C₁₋₆alkyl group, or a C₃₋₆cycloalkyl group;

R¹² is one of:

- (a) a C₃₋₇cycloalkyl group;
- (b) a C₁₋₆alkyl group, optionally substituted with one or more of: a phenyl group; a phenyl group with from 1 to 5 substituents selected from halogen, -NO₂, -CF₃, C₁₋₄alkyl, C₁₋₄alkoxy, -NH₂, -NHCOCH₃, -CONH₂, -OCH₂COOH, -NH(C₁₋₄alkyl), -N(C₁₋₄alkyl)₂,

-62-

-NHCOOC₁₋₄alkyl, -OH, -COOH, -CN and -COOC₁₋₄alkyl; a C₁₋₄alkyl group; a C₁₋₄haloalkyl group; or a halogen; and,

(c) a C₁₋₆perfluoroalkyl group;

5 M represents -S-, -SO-, or -SO₂- ; and,

Het represents a 5 or 6 membered heterocyclic aromatic ring linked to M via a carbon atom of the aromatic ring, said aromatic ring containing 1, 2, 3 or 4 heteroatoms selected from the group consisting of O, N and S said
10 aromatic ring optionally being substituted on carbon atoms of the ring with 1, 2, 3 or 4 substituents selected from the group consisting of -OH, -SH, -CN, -CF₃, NH₂ and halogen.

- 15 2. A compound according to claim 1, wherein:
R¹ and R² are independently -I, -Br, or -Cl.
3. A compound according to claim 1, wherein:
R¹ and R² are both -I.
4. A compound according to any one of claims 1 to 3, wherein:
20 R^{1a}, R^{1b}, R^{2a}, R^{2b} are each independently -H or -CH₃.
5. A compound according to any one of claims 1 to 3, wherein:
R^{1a}, R^{1b}, R^{2a}, R^{2b} are all -H.
- 25 6. A compound according to any one of claims 1 to 5, wherein:
R³ and R⁴ are -CF₃ and -H, respectively.

-63-


7. A compound according to any one of claims 1 to 5,
wherein:
R³ and R⁴ are both -F.
- 5 8. A compound according to any one of claims 1 to 5,
wherein:
R³ and R⁴ are -CF₃ and -H, respectively; and,
R⁵ is -H.
- 10 9. A compound according to any one of claims 1 to 5,
wherein:
R³ and R⁴ are both -F; and,
R⁵ is -F.
- 15 10. A compound according to any one of claims 1 to 5,
wherein:
R³ and R⁴ are both -F; and,
R⁵ is -H.
11. A compound according to any one of claims 1 to 10,
wherein:
Z is -CH₂-T-C(=O)OH or -CH₂-T-C(=O)OR⁸ ; and,
T is -CH₂- .
- 20 12. A compound according to any one of claims 1 to 11,
wherein:
R⁸ is -H, -C(CH₃)₃, or -CH₂-CH=CH₂.
- 25 13. A compound selected from:
{3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoyl}-L-
glutamic acid; and,
the di-tert-butyl ester thereof.

-64-

14. A compound selected from:
3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoyl}-L-glutamic acid; and,
the di-tert-butyl ester thereof.
- 5 15. A compound selected from:
(3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoyl)-L-glutamic acid; and,
the di-tert-butyl ester thereof.
- 10 16. A compound selected from:
(2,3,5-trifluoro-4-[bis(2-chloroethyl)amino]benzoyl)-L-glutamic acid; and,
the di-tert-butyl ester thereof.
- 15 17. A compound selected from:
(2,3,5-trifluoro-4-[bis(2-bromoethyl)amino]benzoyl)-L-glutamic acid; and,
the di-tert-butyl ester thereof.
- 20 18. A compound selected from:
(2,3,5-trifluoro-4-[bis(2-iodoethyl)amino]benzoyl)-L-glutamic acid; and,
the di-tert-butyl ester thereof.
19. A compound selected from:
(3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoyl)-L-glutamic acid; and,
the di-tert-butyl ester thereof.
- 25 20. A compound selected from:
(3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoyl)-L-glutamic acid; and,
the di-tert-butyl ester thereof.

-65-

21. A compound of Formula II:



wherein:

R¹ is -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph ;

5 R² is -Cl, -Br, -I, -OSO₂CH₃, or -OSO₂Ph ;

wherein Ph denotes a phenyl group which is optionally substituted with 1, 2, 3, 4 or 5 substituents independently selected from a C₁₋₄ alkyl group, -F, -Cl, -Br, -I, -CN, or -NO₂;

10 R^{1a} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;

R^{2a} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;

R^{1b} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;

R^{2b} is -H, a C₁₋₄alkyl group, or a C₁₋₄haloalkyl group ;

15 R³ is -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃, -CF₃, -SF₅, -SCF₃, or -CF₂CF₃ ;

R⁴ is -H, -F, -Cl, -Br, -I, -OCHF₂, -C≡CH, -OCF₃, -CH₃, -CF₃, -SF₅, -SCF₃, or -CF₂CF₃ ;

R⁵ is -H or -F;

with the proviso that if R⁴ is -H, then R³ is not -F.

20 22. A compound according to claim 21, wherein:

R¹ and R² are independently -I, -Br, or -Cl.

23. A compound according to claim 21, wherein:

R¹ and R² are both -I.

24. A compound according to any one of claims 21 to 23,

25 wherein:

R^{1a}, R^{1b}, R^{2a}, R^{2b} are each independently -H or -CH₃.

-66-

25. A compound according to any one of claims 21 to 23,
wherein:
 R^{1a} , R^{1b} , R^{2a} , R^{2b} are all -H.
- 5 26. A compound according to any one of claims 21 to 25,
wherein:
 R^3 and R^4 are $-CF_3$ and -H, respectively.
27. A compound according to any one of claims 21 to 25,
wherein:
 R^3 and R^4 are both -F.
- 10 28. A compound according to any one of claims 21 to 25,
wherein:
 R^3 and R^4 are $-CF_3$ and -H, respectively; and,
 R^5 is -H.
- 15 29. A compound according to any one of claims 21 to 25,
wherein:
 R^3 and R^4 are both -F; and,
 R^5 is -F.
- 20 30. A compound according to any one of claims 21 to 25,
wherein:
 R^3 and R^4 are both -F; and,
 R^5 is -H.
31. 3,5-difluoro-4-[bis(2-iodoethyl)amino]benzoic acid.
32. 3,5-difluoro-4-[bis(2-chloroethyl)amino]benzoic acid.
33. 3,5-difluoro-4-[bis(2-bromoethyl)amino]benzoic acid.
- 25 34. 2,3,5-trifluoro-4-[bis(2-chloroethyl)amino]benzoic acid.

-67-

35. 2,3,5-trifluoro-4-[bis(2-bromoethyl)amino]benzoic acid.

36. 2,3,5-trifluoro-4-[bis(2-iodoethyl)amino]benzoic acid.

37. 3,5-difluoro-4-[bis(2-bromopropyl)amino]benzoic acid.

38. 3-trifluoromethyl-4-[bis(2-bromoethyl)amino]benzoic
5 acid.

39. A two-component system comprising:

(i) a first component capable of delivering a
carboxypeptidase enzyme to the interior or exterior
of a target cell or a vector encoding said enzyme
10 to the interior of said cell such that said vector
expresses said enzyme in said cell, and

(ii) a prodrug of according to any one of claims 1 to 20
capable of being converted by said enzyme into a
drug according to any one of claims 21 to 38.

15 40. A kit comprising:

(a) a compound according to any one of claims 1 to 20;
and,

(b) one of:

(i) an immunoglobulin/enzyme fusion protein or
20 conjugate in which the immunoglobulin is specific
for a cellular antigen and the enzyme is a
carboxypeptidase enzyme;

(ii) a ligand/enzyme conjugate or fusion protein,
the ligand being specific for a cellular antigen
25 and the enzyme is a carboxypeptidase enzyme;

(iii) a vector which encodes a carboxypeptidase
enzyme which can be expressed in a cell.

-68-

41. A composition comprising a compound according to any one of claims 1 to 38, and a pharmaceutically acceptable carrier or diluent.
- 5 42. A compound according to any one of claims 1 to 38 for use in a method of treatment of the human or animal body.
43. A compound according to any one of claims 1 to 38 for use in a method of treatment of cancer of the human or animal body.
- 10 44. Use of a compound according to any one of claims 1 to 38 for the manufacture of a medicament for use in the treatment of cancer.
- 15 45. A method for the treatment of cancer comprising administering to a subject suffering from cancer a therapeutically-effective amount of a compound according to any one of claims 1 to 38.

1/1

Figure 1A





Figure 1B



INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/01194

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C237/36 A61K31/223 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 25429 A (CANCER RES CAMPAIGN TECH ; SPRINGER CAROLINE JOY (GB)) 10 November 1994 (1994-11-10) page 22	1,2,4,5, 11,12, 41-45
X	WO 97 03957 A (PHARMACIA & UPJOHN SPA ; COZZI PAOLO (IT); BERIA ITALO (IT); CAPOLO) 6 February 1997 (1997-02-06) page 14	21,22, 24,25

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

25 July 2000

Date of mailing of the international search report

02/08/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 eponl,
Fax: (+31-70) 340-3016

Authorized officer

Bader, K

INTERNATIONAL SEARCH REPORT

information on patent family members

Inter national Application No

PCT/GB 00/01194

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9425429 A	10-11-1994	AT 172958 T	15-11-1998
		DE 69414379 D	10-12-1998
		DE 69414379 T	12-05-1999
		EP 0696270 A	14-02-1996
		ES 2125450 T	01-03-1999
		JP 8509490 T	08-10-1996
		US 5811454 A	22-09-1998
WO 9703957 A	06-02-1997	AU 6357996 A	18-02-1997
		BR 9606528 A	23-12-1997
		CA 2199635 A	06-02-1997
		CN 1159183 A	10-09-1997
		EP 0787126 A	06-08-1997
		HU 9702393 A	28-04-1998
		JP 10506410 T	23-06-1998
		NO 971142 A	12-03-1997
		PL 319352 A	04-08-1997

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☒ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.